

BIOCONTROL



ISOMATE®-C PLUS

ACTIVE INGREDIENTS:

(E, E)-8, 10-Dodecadien-1-ol 53.0%
Dodecanol 29.7%
Tetradecanol 6.0%

OTHER INGREDIENTS 11.3%

TOTAL 100.0%

182.3 mg active ingredients per dispenser

Keep out of reach of children
WARNING

FIRST AID

If on Skin or Clothing * Take off contaminated clothing.
* Rinse skin immediately with plenty of water for 15-20 minutes.
* Call a poison control center or doctor for treatment advice.

If in Eyes * Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
* Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-999-8805 for further questions.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear eye protection (such as goggles or safety glasses) and gloves when handling. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of this product. Do not discharge into lakes, streams, ponds or public waters unless in accordance with NPDES permit. For guidance, contact your regional office of the Environmental Protection Agency.

PACIFIC BIOCONTROL CORPORATION

14615 NE 13th Court, Suite A, Vancouver, WA 98685 U.S.A.
Telephone (360) 571-2247 or 1-800-999-8805

MADE IN JAPAN
EPA Est. No: 47265-JP-01
EPA Reg. No: 53575-6

NET CONTENTS: 400 Dispensers Units
One dispenser contains 0.008 fl oz or 205.4 mg
Total content of package: 3.2 fl oz or 82.2 gm

ISOMATE® is a registered Trademark of Pacific Biocontrol Corporation

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

Pesticide Storage Store unopened package at temperatures below 40°F in a dry location. Product may be stored in cold storage facilities used for food storage.

Pesticide Disposal Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal Dispose of dispensers and envelopes in sanitary landfill or by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

It is critical that ISOMATE-C PLUS is applied as directed.

- 1. Crop Apple, pear, walnut, quince, prune, plum, peach, pecan, and nectarine.
2. Pest Codling moth (Cydia pomonella), hickory shuckworm (Cydia caryana).
3. Rate 400 dispensers per acre or 1000 dispensers per hectare (2.9 fl oz or 72.9 gm a.i. per application). Apply double rate of dispensers to edges of orchard. Do not exceed 150 gm a.i. per acre per year.

4. Application Apply dispensers securely to lateral branches in upper third of tree canopy, preferably within 2-3 feet of treetop. Dispensers twisted too tightly may girdle branches. Can be applied efficiently from a moving trailer or with a pole applicator.

5. Timing Apply prior to moth emergence in spring. Dispensers release pheromone for up to 120-140 days depending on temperature. In crops with long field seasons (i.e., more than 120 days), a second application is recommended. If subsequent applications are required, apply prior to the start of subsequent flights. Consult your local pest control advisor for proper timing.

6. Note Isomate-C Plus suppresses codling moth and hickory shuckworm from mating. However, if a major source of mated female moths of these species is present adjacent to the treated orchard, migration of these moths may significantly reduce the level of control achieved. Sources are likely to be unsprayed apple, pear, quince, walnut, plum, crabapple trees, or other host species within 300 yards of the treated orchard. This can be overcome by:
a. Treatment of entire blocks and not just sections of large conventionally treated orchards that frequently serve as sources of mated females.
b. Treatment of infestation sources with Isomate-C Plus (i.e., a strip at least 50 yards wide nearest the treated orchard).
c. Treatment of infestation source with an effective insecticide.
d. Treatment of 4-6 rows along border or pheromone treated orchard with insecticide. Supplementary applications of insecticide are advised when Isomate-C Plus is used to control high populations of codling moth or hickory shuckworm. All pests must be monitored on a regular schedule so that timely intervention with insecticides is possible.

WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed. The Manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

Revised Date: December 2004