

BIOCONTROL



ISOMATE®-CM/LR TT

**A MATING DISRUPTION FORMULATION FOR CODLING MOTH (*Cydia pomonella*),
OBLIQUEBANDED LEAFROLLER (*Choristoneura rosaceana*),
PANDEMIS LEAFROLLER (*Pandemis pyrusana*), FRUITTREE LEAFROLLER
(*Archips argyrospilus*), THREELINED LEAFROLLER (*Pandemis limitata*),
EUROPEAN LEAFROLLER (*Archips rosanus*).**

ACTIVE INGREDIENTS:

(E, E)-8, 10-Dodecadien-1-ol	38.62%
1-Dodecanol	6.00%
1-Tetradecanol	1.40%
Z-11-Tetradecen-1-yl Acetate	38.04%
Z-9-Tetradecen-1-yl Acetate	4.34%
Z-11-Tetradecen-1-ol	1.05%
Z-11-Tetradecenal	1.00%

OTHER INGREDIENTS 9.55%

TOTAL 100.0%

328.13 mg active ingredients per dispenser

Keep out of reach of children

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes, or clothing. Harmful if absorbed through skin. Causes moderate eye irritation. After handling, wash thoroughly with soap and water before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before reuse. Wear chemical resistant gloves and protective eyewear.

FIRST AID STATEMENT

IF ON SKIN	* Remove contaminated clothing.
OR	* Immediately rinse skin for 15-20 minutes using plenty of water.
CLOTHING	* Contact a poison control center or doctor for treatment advice.
IF IN EYES	* Immediately hold eyes open, slowly and gently rinse with water for 15-20 minutes. If wearing contact lenses, rinse eyes for 5 minutes, remove contact lenses, and continue rinsing. * Contact a poison control center or doctor for treatment advice.

HOT LINE NUMBER: Have the product container or label with you when contacting a poison control center or doctor, or going for treatment. For emergency information on this product, call the **National Pesticides Information Center** AT 1-800-858-7878, 6:30AM to 4:30PM Pacific time (PT), seven days a week. During other times, call the poison control center at 1-800-222-1222. You may also contact Pacific Biocontrol at 1-800-999-8805 for additional information.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of this product.

PACIFIC BIOCONTROL CORPORATION

14615 NE 13th Court, Suite A, Vancouver, WA 98685 U.S.A.
Telephone (360) 571-2247 or 1-800-999-8805

MADE IN JAPAN

EPA Est. No: 47265-JP-01

EPA Reg. No: 53575-31

NET CONTENTS: 300 Dispensers Units

One dispenser contains 0.014 fl oz or 362.77 mg

Total content of package: 4.22 fl oz or 108.83 gm

ISOMATE® is a registered Trademark of Pacific Biocontrol Corporation

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

It is CRITICAL that ISOMATE-CM/LR TT is applied as directed.

Crops	Apples, pears, quince and other pome fruits; peaches, prunes, plums, nectarines, cherries and other stone fruits; walnut, pecan and other tree nut crops.
Target Pests	Codling moth (<i>Cydia pomonella</i>), obliquebanded leafroller (<i>Choristoneura rosaceana</i>), pandemis leafroller (<i>Pandemis pyrusana</i>), fruittree leafroller (<i>Archips argyrospilus</i>), threelined leafroller (<i>Pandemis limitata</i>), European leafroller (<i>Archips rosanus</i>).
Rate	300 dispensers per acre (3.82 fl oz or 98.44 gm a.i. per application). Apply double rate of dispensers to edges of orchard. Do not exceed 150 gm a.i. (or 457 dispensers) per acre per year.
Application	Place dispensers on lateral branches in upper third of tree canopy. Apply dispensers uniformly in the treated area.
Timing	Apply prior to codling moth emergence in spring. Consult your local pest control advisor or Pacific Biocontrol representative for proper timing or estimated dispenser longevity in your area.
Note	Isomate-CM/LR TT suppresses mating of target pests. Immigration of mated female moths of these species from adjacent external orchards of infestation will reduce the level of control. Manage by: a. Treatment of external sources of infestation with Isomate-CM/LR TT. b. Treatment of external sources of infestation with an effective insecticide. c. Treatment of pheromone treated orchard with insecticide. Areawide application is most effective. Supplementary applications of insecticide are advised when Isomate-CM/LR TT is used in orchards with high pest populations. All pests must be monitored so that timely intervention with insecticides is possible.
STORAGE AND DISPOSAL	
Do not contaminate water, food or feed by storage or disposal	
Pesticide Storage	Store in original unopened package in a dry location at temperatures below 40°F. Only unopened or unbroken dispenser packages may be stored in cold facilities used for food storage.
Pesticide Disposal	Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.
Container Disposal	Nonrefillable container. Do not reuse or refill this container. Dispose of empty foil packet in the trash.

WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed. Pacific Biocontrol Corporation warrants that the product complies with the specifications expressed in this label. To the extent consistent with applicable law, Pacific Biocontrol Corporation makes no other warranties, and disclaims all other warranties, expressed or implied, including but not limited to warranties or merchantability and fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. To the extent consistent with applicable law, Pacific Biocontrol Corporation's liability or default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Pacific Biocontrol Corporation shall have no liability for damages (special, consequential or incidental) that result from handling, storage and use of this product, which is not in compliance with the label. Use of this product implies all directions of this label have been read carefully. Read all directions carefully.

ISOMATE® is a registered Trademark of Pacific Biocontrol Corporation

Revision Date: March 2010